

Echokardiográfiás alapismeretek

Kardiomiopátiák

Dr. Habon Tamás
PTE KK I.sz. Belgyógyászat
Kardiológiai Tanszék

ACCF/ASE/AHA/ASNC/HFSA/HRS/SCAI/SCCM/SCCT/SCMR

2011 Appropriate Use Criteria for Echocardiography

A Report of the American College of Cardiology Foundation Appropriate Use Criteria Task Force, American Society of Echocardiography, American Heart Association, American Society of Nuclear Cardiology, Heart Failure Society of America, Heart Rhythm Society, Society for Cardiovascular Angiography and Interventions, Society of Critical Care Medicine, Society of Cardiovascular Computed Tomography, and Society for Cardiovascular Magnetic Resonance

Endorsed by the American College of Chest Physicians


Cardiomyopathies With TTE

• Initial evaluation of known or suspected cardiomyopathy (e.g., restrictive, infiltrative, dilated, hypertrophic, or genetic cardiomyopathy)	A (9)
• Re-evaluation of known cardiomyopathy with a change in clinical status or cardiac exam or to guide therapy	A (9)
• Routine surveillance (<1 y) of known cardiomyopathy without a change in clinical status or cardiac exam	I (2)
• Routine surveillance (≥1 y) of known cardiomyopathy without a change in clinical status or cardiac exam	U (5)
• Screening evaluation for structure and function in first-degree relatives of a patient with an inherited cardiomyopathy	A (9)
• Baseline and serial re-evaluations in a patient undergoing therapy with cardiotoxic agents	A (9)

Ischemic Cardiomyopathy/Assessment of Viability With Stress Echocardiography

• Known moderate or severe LV dysfunction • Patient eligible for revascularization • Use of dobutamine stress only	A (8)
--	-------


PRIMER CARDIOMYOPATHIÁK
(elsődlegesen a szívet érintik)



<http://www.genedx.com>

Maron BJ et al; Circulation 2006;113:1807-1816

Cardiomyopathiák



Classification of the cardiomyopathies: a position statement from the European Society Of Cardiology Working Group on Myocardial and Pericardial Diseases. **Eur Heart J. 2008; 29:270-276.**


CM: Specifikus etiológiák

- Ischemic
- Valvular
- Hypertensive
- Inflammatory
- Metabolic
- Inherited
- Toxic reactions
- Peripartum

Ischemic: thinned, scarred tissue



DCM



DCM echocardiográfiás jellemzői

- Bal kamrai dimenziók
 - Átmérő
 - Spercitás
- Bal kamra systoles funkciója
 - EF
 - CO
 - dP/dt
- Segmentális falmozgászavarok
 - Etiológia??
 - Stress echo
- Bal kamra diastoles funkciója
 - E/A
 - E/E'
- Mitrális regurgitáció
 - Funkcionális
 - Ischemiás
- Bal kamrai thrombus
- Jobb kamra funkció
 - TAPSE
- Pulmonális nyomás
 - IVC
 - TR

Hogyan mérjük a bal kamrai átmérőket?

- Ínhúrok szintjében, merőlegesen az elképzelt hossz tengelyre


■ 2D irányított M-mód

- parasternális hossz tengely
 - Falak merőlegesen a sugárirányra
- Esetleg parasternális rövid tengely
- Jobb időbeli felbontás (magasabb frame rate)
- Gyakran „off axis”

■ 2D


- Parasternális hossz tengely
- Csúcsról lehetőleg ne!
- Rosszabb időbeli felbontás
- Minden esetben merőleges a képzeletbeli hossz tengelyre

Átmérők: 2D irányított M-mód





- Végdiastole
 - QRS kezdete

- Végysystole
 - Legkisebb átmérő
 - Aszinkronia esetén a posterior fal mozgása számít



Sphericity index

- Normálisan a hosszátmérő és rövidebb átmérő aránya > 1.6 .
- DCM-ben < 1.5 , akár egyenlő is lehet
- Prognosztikus szerepe van







Ejekciós frakció: Simpson módszer


- Biplane módszer:
 - Az egymásra kb. merőleges csúcsi 4,-ill. 2-üregi metszetben mérünk areát
- Csak jól látható vér-endocardium határ esetén pontos!
- Papilláris izmot és trabeculákat volumenként rajzoljuk
- 3D!!! Kontraszt

EDV

ESV



Contrastecho




L

HR : 57



15/09/2012 13:05:32
 Freq.: 5.0 MHz/5.0 MHz
 FPS: 24.9

EDV	175 ml
ESV	82 ml
EF	53 %
HR	55 BPM
SV	93 ml
CO	5.1 l/min
Spl	0.35



CO: hogyan mérjük?

- LVOT átmérő mérése systoleban, parasternális hossztengetlyi nézetben
- Átmérőből LVOT areát számolunk
- D^2 : a mérési hibát is négyzetre emelem!
- VTI: távolság jellegű paraméter
- Csúcsi ötüregű nézetből az LVOT-ban, pulzatis Dopplerrel!
- 3 (5) mérést átlagolunk






$$SV = VTI \times A$$

Globális systoles functio: dP/dt hogyan mérjük?



- MI sebesség spectrum CW-vel mérve
- Mennyi idő telik el az 1 m/s és a 3 m/s sebesség elérése között (32 Hgmm-nyi nyomásemelkedés)

$$dP/dt = \frac{32}{\Delta t}$$


- Hosszabb Δt ill. alacsonyabb dP/dt (< 600 Hgmm/sec) károsodott BKF-ra utal




Diastolés funkció megítélése



Normális **Károsodott relaxáció** **Pseudonormális** **Restriktív**




- A mitrális beáramlási görbe preload-függő, a normális és pseudonormális görbe nem megkülönböztethető






- Klasszikusan a véna pulmonális beáramlási görbe segít a feladat megoldásában

Diasztolés funkció megítélése





E/E' szerepe





- Az emelkedett E/E' emelkedett bal kamrai töltőnyomásra utal. Segíthet a normális és pseudonormális mitrális beáramlási görbe elkülönítésében

Non-ischemiás DCM: funkcionális mitrális regurgitáció




- A gömbszerű kamrában a papilláris izmok apicalis, laterális irányba tolódnak el, ami a mitrális felfüggesztő apparátus megrövidülését eredményezi




- A vitorlák koaptációja romlik, ami többé-kevésbé centrális regurgitációs jet-et eredményez

Ischemiás eredetű mitrális regurgitáció

- Ischemiás szívben az egyik mitrális vitorla hegesedik, mozgásában korlátozottá válik
- Gyakrabban a posterior vitorlát érinti inferior infarctus vagy ischemia talaján
- A mellső vitorla prolapsusát utánozza, excentrikus jet-et eredményez
- Itt nem „az ép alá fúj”!




Mitral regurgitation




Bal kamrai thrombus

- Ischemiás DCM-ben, szegmentális falmozgászavar esetén a bal kamrai thrombus gyakoribb
- Non-ischemiás DCM-ben, diffúz hypokinesis esetén is előfordul





- A spontán echokontraszt pre-thromboticus állapot, de thrombus nem minden esetben van jelen

DCM - SEC



Bal kamrai thrombus

- Ischemiás DCM-ben, szegmentális falmozgászavar esetén a bal kamrai thrombus gyakoribb
- Non-ischemiás DCM-ben, diffúz hypokinesis esetén is előfordul





Jobb kamra funkció

- Tricuspid annular plane systolic excursion (**TAPSE**)
 - Kóros < 24 mm
 - < 14 mm igen rossz prognózist jelent




- A jobb kamra funkció károsodása rosszabb prognózist jelent





Jobb kamra nehézségei

2D, 3D, TAPSE, Doppler

Előről nézve
háromszög alakú






Keresztmetszetben
félhold alakú



Pulmonális nyomás

- Emelkedése rosszabb prognózist jelent
- A tricuspidalis insuffitientia sebességéből indirekt módon kiszámolható a jobb kamrai systolés nyomás
 - Systolés PAP= $4v^2 + \text{RA nyomás}$
- Több nézetből nézzük meg
- Pulmonalis stenosis vagy RVOT obstructio jelenlétében nem reális





Jobb pitvari nyomás becslése

- IVC tágassága ill. az átmérő változása belégzésben



IVC size	Collapsibility Index	RA Pressure
Normal IVC size (≤ 1.7 cm)	50% decrease in size	0 - 5 mm Hg
Dilated IVC (>1.7 cm)	$\geq 50\%$ inspiratory collapse	6 – 10 mm Hg
	$< 50\%$ inspiratory collapse	10 – 15 mm Hg
	no collapse	> 15 mm Hg

Source: Lang RM, Bierig M, Devereux RB, et al. Journal of the American Society of Echocardiography 18:1458, 2005.



„Free” tricuspidális regurgitáció

- Erősen dilatált jobb kamra, vagy a tricuspidális billentyű organikus betegsége esetén a jobb kamra és pitvar közötti nyomáskülönbség kicsi



- A tricuspidális regurgitáció sebessége is alacsony. Az ebből kalkulált nyomás nem tükrözi a pulmonális nyomást

CRT




Figure 4. Quantification of regional wall motion from real-time three-dimensional echocardiographic data. After semiautomatic segmentation of the left ventricular chamber (**upper left**), the extent and timing of regional wall motion is analyzed in a 16-segment model (**lower left**) and in this example expressed as regional ejection fraction over time. There is clear regional dyssynchrony between the inferoseptal and the anterolateral segments during left bundle branch block (LBBB) (**lower middle**), which improves immediately after initiation of cardiac resynchronization therapy (CRT) (**lower right**). (Courtesy of A. Franke, University Hospital Aachen, Germany.)








FIGURE 14–30B A, Diagrammatic cross sections of the left ventricle to show 17 segments for wall motion analysis. **B**, Left ventricular segments from three apical views and mid parasternal short-axis view with the corresponding coronary artery distribution.


Cx = circumflex artery; LAD = left anterior descending artery; RCA = right coronary artery.

(Modified from Lang RM, Bierig M, Devereux RB, et al, Chamber Quantification Writing Group; American Society of Echocardiography's Guidelines and Standards Committee; European Association of Echocardiography: Recommendations for chamber quantification: A report from the American Society of Echocardiography's Guidelines and Standards Committee and the Chamber Quantification Writing Group, developed in conjunction with the European Association of Echocardiography, a branch of the European Society of Cardiology. *J Am Soc Echocardiogr* 18:1440, 2005. Used with permission.)



Hibrid képalkotás

Stress MPS + MSCT



HYPERTROPHIAS CARDIOMYOPATHIA (HCM)



Myocardial Hypertrophy:
Primarily septal

Myocardial Fiber Disarray
Interstitial Fibrosis

**Teare : Asymmetrical hypertrophy of the heart in young adults.
Br Heart J 1958.**

Echo vizsgálati lehetőségek HCM-ben





- Hagyományos módszerek (2D, M-mód, Doppler)
- Szöveti Doppler, Color M-mód flow propagáció, strain, speckle tracking-BK torzió analízis, kontrasztechó, 3D echo
 - Falvastagság, izomtömeg
 - Lokalizáció
 - SAM, LVOT obstructio,
 - Szisztolés f., Diasztolés f.
 - Bal pitvar area, volume,
 - Dyssynchronia (diasztolés is!)



Echocardiographia szerepe HCM-ben

- Hagyományos módszerek (2D, M-mód, Doppler)
- Szöveti Doppler, Color M-mód flow propagáció, strain, speckle tracking-BK torzió analízis, kontrasztechó, 3D echo
 - Falvastagság, izomtömeg
 - RV hypertrophia?, RV obstrukció?
 - LV-EF
 - LV-Diasztolés funkció
 - A. pulmonális syst. nyomás
 - Bal pitvar area, volumen, index
 - LVOT dinamikus obstrukció, SAM
 - Mitrális billentyű, papilláris izom
 - Myectomy, alkoholos abláció (TTE/TEE)

HCM-falvastagság

- Elsősorban septum basalis része érintett
- Egyéb lokalizáció:
 - Koncentrikus
 - Apicális
 - Lateralis fal....
 - Jobb kamra
- Falvastagság meghatározása PSAX (basalis, mid-LV, apex)
 - Maximális falvastagság
 - Hypertrophia kiterjedése
 - PLAX: túlmérés veszélye

A**B****C****D**




84
2:52 HR

Courtesy of the AMC Echolab, AMC, The Netherlands

ECHO PEDIA .ORG
portal of cardiac ultrasound

09/11/2009 14:51:29



Mitrális mellső vitorla SAM jelensége




■ SAM

- nincs
- nyugalmi
- terhelésre jelentkezik csak

■ Következmények


- LVOTO (>30 Hgmm)
- MR

Pathophysiology of LVOT Obstruction and MR: Grigg et al JACC 1993




Venturi-hatás







A TDI alkalmas a pathológiás és physiologiás bal kamra hypertrophia elkülönítésére






• Szöveti Doppler görbe a laterális anuluson mérve (a. HCM, b. hypertonia, c. egészséges személy, d. atléta)

- A longitudinális funkciót nem a bal kamra hypertrophia mértéke, hanem pathológiás volta (fibrózis) befolyásolja
- HCM-ben és hypertoniás szívben S és Ea egyaránt károsodott, míg atlétákban közel normális



Apical HCM








FIGURE 14–17 Apical view showing a mid cavity obstruction (arrows) that was not visualized clearly with 2D echocardiography without contrast injection. An apical pouch (A) is clearly seen.


(From Oh JK, Seward JB, Tajik AJ: The Echo Manual. 3rd ed. Philadelphia, Lippincott Williams & Wilkins, 2006. Used with permission of Mayo Foundation for Medical Education and Research.)


Intracoronary myocardial contrast echocardiography in the harmonic imaging mode demonstrated that the patient was unsuitable for alcohol septal ablation.



La Canna G *et al.* (2007) Alcohol septal ablation versus surgical myectomy: a patient with obstructive HCM
Nat Clin Pract Cardiovasc Med 4: 570–576 doi:10.1038/ncpcardio0988










- 2-The presence of LVH is not the end, but the beginning of the story
- 3- The correct diagnosis is important in therapy & prognosis

Table 5 Echocardiographic features that suggest specific aetiologies (modified from Rapezzi et al.⁶⁷)

Finding	Specific diseases to be considered
Increased interatrial septum thickness	Amyloidosis
Increased AV valve thickness	Amyloidosis; Anderson-Fabry disease
Increased RV free wall thickness	Amyloidosis, myocarditis, Anderson-Fabry disease, Noonan syndrome and related disorders
Mild to moderate pericardial effusion	Amyloidosis, myocarditis
Ground-glass appearance of ventricular myocardium on 2D echocardiography	Amyloidosis
Concentric LVH	Glycogen storage disease, Anderson-Fabry disease, PRKAG2 mutations
Extreme concentric LVH (wall thickness ≥ 30 mm)	Danon disease, Pompe disease
Global LV hypokinesia (with or without LV dilatation)	Mitochondrial disease, TTR-related amyloidosis, PRKAG2 mutations, Danon disease, myocarditis, advanced sarcomeric HCM, Anderson-Fabry disease
Right ventricular outflow tract obstruction	Noonan syndrome and associated disorders

2014 ESC Guidelines on diagnosis and management of hypertrophic cardiomyopathy: the Task Force for the Diagnosis and Management of Hypertrophic Cardiomyopathy of the European Society of Cardiology (ESC). Eur Heart J. 35(39), 2733-79, 2014.

2D 3D AND 3LAYER STE HOM



Specific patterns of L,R, C, rotation, twist & torsion mechanics

reduced systolic strain (L,R,C), reduced apical systolic and diastolic rotation rates, delayed untwist

Useful in the differential diagnosis????

Conventional Diagnostic Criteria for HCM

Non Apical HCM variants: IVS/PW: >1.3, IVS >15mm, SAM/ LVOT obstruction

Apical HCM: exuberant apical LVH, spade-like LV cavity

HCM Diagnosis Unequivocal

Consider Risk Stratification

Echo-Doppler Predictors of Adverse Prognosis Non-Apical variants

- LVH (>30 mm)
- LVOT gradient (>30mm Hg)
- LV dysfunction (EF < 50 %)
- LA dilation (> 48 mm)^{*} or LAVI >34 ml/m²
- Intraventricular dyssynchrony (>45 ms)^{*} ^{*}

HCM-Diagnosis Equivocal

Non-Apical HCM variants

Differentiation from Pathologic LVH Criteria favoring HCM

- TDI longitudinal strain(> -10 %)
- Paradoxical longitudinal strain linear mapping-TDI)
- Attenuated regional radial and circumferential strain(2D strain)
- Intraventricular dyssynchrony

Apical HCM

- 2 D contrast-enhanced echo
- RT-3DE
- RT-3DE with contrast
- 2 D-strain (paradoxical apical strain)

Differentiation from Physiologic LVH and Normal[†] Criteria favoring HCM

- Systolic mitral annular velocity(< 9 cm/s)
- Diastolic dysfunction (TDI)
- Lack of LVH regression post cessation of exercise
- Lack of LV dilation
- Provocable LVOT gradient
- Attenuated of longitudinal, circumferential and radial strain (2 D strain)[†]
- Intraventricular Dyssynchrony[†]

RCM - OSZTÁLYOZÁS

MYOCARDIALIS

- Noninfiltratív - idiopathiás, scleroderma, diabetes
- Infiltratív - amiloid, sarcoid, Gaucher kór
- Tárolási betegségek - haemochromatosis, Fabry kór, stb.

ENDOMYOCARDIALIS



- Endomyocardialis fibrosis (EMF)
- Hypereosinophil syndrome (HES)
- Carcinoid
- Metastaticus malignómák
- Irradiáció
- Anthracycline toxicitás

RCM echocardiographiás jellemzői

- Kisebb vagy normális balkamrai üregméretek
- Balkamra hypertrophia nincs/enyhe
- Igen tág pitvarok, tág IVC
- Jó/enyhén csökkent systoles BKF
- Súlyos diasztoles funkciózavar



Amyloid szív

- ECHO-val vastag falak, csillámlás; EKG-n low voltage



- Károsodott systolés és diasztolés longitudinális funkció


Amyloid szív



Amyloid Heart Disease – Echo Features

- Ground glass pattern
- Left ventricular hypertrophy
- Atrial enlargement
- Thickened interatrial septum
- Thickened valves frequently with mild regurgitations
- Advanced diastolic dysfunction
- Pericardial/Pleural effusion
- “Apical sparing pattern” of longitudinal strain
- Systolic dysfunction (endstage)
- Right heart involvement


The echocardiogram is often so typical that it leads to the diagnosis of amyloidosis.



AMYLOIDOSIS – apical four-chamber view/2D

Typical features of amyloidosis, including echogenic/hourglass appearance of the myocardium, thickened valves, and enlarged atria. This patient also received a pacemaker.

30/03/2016 11:42:00



21/07/2015 07:16:25

V


5

X


10

15

100
HR




21/07/2015 07:17:27



Endomyocardialis Fibrosis

Endemic in parts of Africa, India, South and Central America, Asia
15-25% of cardiac deaths in equatorial Africa
hypereosinophilic syndrome (Löffler's endocarditis)



Thickening of basal inferior wall
endocardial deposition of thrombus
apical obliteration
mitral regurgitation
80-90% die within 1-2 years

Some authors suggest that the binary sign, defined as binary appearance of the left ventricular endocardial border, aids in the diagnosis of Fabry's disease.


However, the sensitivity and specificity of this sign is rather low.

FABRY'S DISEASE – apical four-chamber view/2D

Pronounced bi-ventricular hypertrophy and rather speckled appearance of the myocardium.

Fabry's Disease: Echo Features


- Left ventricular hypertrophy
- Right ventricular hypertrophy
- Myocardial fibrosis
- Diastolic dysfunction/enlarged left atria



Sarcoid Heart Disease – Echo Features

- Cardiac involvement in sarcoidosis is associated with a poor prognosis
- Pericardial effusion
- Left ventricular aneurysms
- Wall motion abnormalities (not related to coronary perfusion territories)
- Hypertrophy (segmental)
- Edema/Fibrosis
- End-stage: left ventricular dilatation, wall thinning and impaired left ventricular function


20 – 30 % of patients with proven sarcoidosis have cardiac involvement. MRI is more sensitive than echo in the detection of sarcoid heart disease.



SARCOIDOSIS – apical four-chamber view/2D


Abnormal cardiac geometry with segmental wall motion abnormalities, thickening, and increased echogenicity in the region of the mid- and distal anterior septum.

Carcinoid heart disease.



Scarsbrook A F et al. Radiographics 2007;27:455-477

RadioGraphics



75
3:87 HR

Courtesy of the AMC Echolab, AMC, The Netherlands

ECHO PEDIA.ORG
part of cardiac ultrasound.org



Mitrális “A” hullám és PV “a” hullám

■ Restrictive Physiology:


- PV-a Velocity > 35 cm/s

OR

- PV-a duration, 30 ms longer than Mitral “A” wave duration.



Hepatic Vein Dopplar: Normal




Systolic and diastolic forward flow

S-vel. > D-vel.

Diastolic flow reversal:
Expir. >> Insp.


Máj véna Doppler: Restriction



Forward flow primarily in Diastole.



Inspiration increases both
>systolic, and
>Diastolic
Flow reversals.

Constrictiv pericarditis




Constrictiv vagy restrictiv folyamat?

Szöveti Doppler!



- Az anuluson mért kora diasztolés sebesség restrictiv cardiomyopathiában csökkent, míg pericardiális constrictioban normális


Májvéna Doppler: Constriction



Diastolic flow reversal is augmented in expiration.


$DFR_{exp.} > 25\%$ forward diastolic velocity

Constriction Doppler










Pericardialis tamponade



Tamponade - Paradox pulsus




ARVD - MRI



1:5000
Genetika (desmosoma)

ARVD 2D echo eltérések




Echocardiographic views from probands meeting Task Force Criteria for arrhythmogenic right ventricular (RV) dysplasia



Yoerger, D. M. et al. J Am Coll Cardiol 2005;45:860-865

- segmental RWMA
- focal RV aneurysms
- RVOT/RVIT dilatation
- RV systolic dysfunction
- trabecular derangement
- moderator band hyper-reflectivity

ARVD



Global or regional dysfunction or structural alterations

Modified Task Force Criteria

Major (By 2D echo)

- Regional RV akinesia, dyskinesia or aneurysm
and 1 of the following (end diastole)
 - ✓ PLAX RVOT $\geq 32\text{mm}$ ($\geq 19\text{mm/m}^2$)
 - ✓ PSAX RVOT $\geq 36\text{mm}$ ($\geq 21\text{mm/m}^2$)
 - ✓ or fractional area change $\leq 33\%$

Minor (By 2D echo)

- Regional RV akinesia, dyskinesia or aneurysm
and 1 of the following (end diastole)
 - ✓ PLAX RVOT $\geq 29\text{mm}$ to $< 32\text{mm}$ ($\geq 16\text{mm/m}^2$ to 19mm/m^2)
 - ✓ PLAX RVOT $\geq 32\text{mm}$ to $< 36\text{mm}$ ($\geq 18\text{mm/m}^2$ to 21mm/m^2)
 - ✓ or FAC $> 33\%$ to $\leq 40\%$

Original Task Force Criteria


Major

- Severe dilatation and reduction of RV EF with no or mild LV involvement
- Localized RV aneurysms (akinetic or dyskinetic areas with diastolic bulging)
- Severe segmental dilatation of the RV

Minor

- Mild global RV dilatation and/or EF reduction with normal LV
- Mild segmental dilatation of the RV
- Regional RV hypokinesia

NEM OSZTÁLYOZHATÓ CMP





■ Left ventricular non-compaction

- Barth sy. - Tafazzin (X linked)
- Lamin A/C, ZASP, alpha-dystrobrevin
- Echocardiographia (Chin 1990, Jenni 2001)

■ Tako Tsubo cardiomyopathy

- Magas epinephrin szint pathophysiologiai szerepe
- Lokalizáció, MR, Biventricularis





75
3:46 HR

Courtesy of the AMC Echolab, AMC, The Netherlands


ECHO PEDIA.ORG
part of cardiac ultrasound.org

NON COMPACT CMP

Chin 1990




1. multiple trabeculations
2. two-layer structure
3. X to Y ratio < 0.5
4. SA and apical views



Jenni 2001



1. multiple trabeculations
2. deep recesses
3. two-layer structure
4. NC / C ratio > 2
5. Parasternal SA view



Stöllberger 2002




1. > 3 trabeculations
2. Apically to the PM
3. In 1 image plane
4. Intertrabecular spaces
5. Apical views




Speckle tracking

Medscape




Source: Heart © 2013 BMJ Publishing Group Ltd & British Cardiovascular Society

TAKO-TSUBO (STRESS) CMP



DIASTOLE



SYSTOLE

Köszönöm a figyelmet!

